

## REFERENCES

1. Cruz PD, Nixon-Fulton J, Tigelaar RE, Bergstresser PR: Disparate effects of in vitro low-dose UVB irradiation on intravenous immunization with purified epidermal cell subpopulations for the induction of contact hypersensitivity. *J Invest Dermatol* 92:160-165, 1989
2. Cooper KD, Fox P, Neises G, Katz SI: Effects of ultraviolet radiation on human epidermal cell alloantigen presentation: Initial depression of Langerhans cell-dependent function is followed by the appearance of T6-DR+ cells that enhance epidermal alloantigen presentation. *J Immunol* 134:129-137, 1985
3. Baadsgaard O, Wulf HC, Wantzin GL, Cooper KD: UVB and UVC, but not UVA, potently induce the appearance of T6-DR+ antigen-presenting cells in human epidermis. *J Invest Dermatol* 89:113-118, 1987
4. Baadsgaard O, Fox DA, Cooper KD: Human epidermal cells from ultraviolet light-exposed skin preferentially activate autoreactive CD4+ 2H4+ suppressor-inducer lymphocytes and CD8+ suppressor/cytotoxic lymphocytes. *J Immunol* 140:1738-1744, 1988
5. Takeuchi T, Schlossman SF, Morimoto C: The 2H4 molecule but not the T3-receptor complex is involved in suppressor inducer signals in the AMLR system. *Cell Immunol* 107:107-114, 1987
6. Baadsgaard O, Mannie A, Dass B, Fox DA, Cooper KD: Increased T-suppressor cell activity in humans after stimulation by UV-exposed epidermal cells: Involvement of the 2H4 molecule. *J Invest Dermatol* 90:544, 1988
7. Baadsgaard O, Lisby S, Wantzin GL, Wulf HC, Cooper KD: Rapid recovery of Langerhans cell alloreactivity, without induction of autoreactivity, following in vivo UVA, but on UVB exposure of human skin. *J Immunol* 142:4213-4218, 1989

## REPLY

We thank Dr. Baadsgaard for taking the time to correspond and rectify the references indicated.

Ponciano D. Cruz, Jr., MD  
Department of Dermatology  
University of Texas  
Southwestern Medical Center at Dallas  
Dallas, Texas